

Culture and PCR Detection of Haemophilus influenzae and Haemophilus haemolyticus in Australian Indigenous Children with Bronchiectasis

K. M. Hare, M. J. Binks, K. Grimwood, A. B. Chang, A. J. Leach and H. Smith-Vaughan
J. Clin. Microbiol. 2012, 50(7):2444. DOI:
10.1128/JCM.00566-12.
Published Ahead of Print 2 May 2012.

Updated information and services can be found at:
<http://jcm.asm.org/content/50/7/2444>

	<i>These include:</i>
REFERENCES	This article cites 12 articles, 4 of which can be accessed free at: http://jcm.asm.org/content/50/7/2444#ref-list-1
CONTENT ALERTS	Receive: RSS Feeds, eTOCs, free email alerts (when new articles cite this article), more»

Information about commercial reprint orders: <http://journals.asm.org/site/misc/reprints.xhtml>
To subscribe to to another ASM Journal go to: <http://journals.asm.org/site/subscriptions/>

Culture and PCR Detection of *Haemophilus influenzae* and *Haemophilus haemolyticus* in Australian Indigenous Children with Bronchiectasis

K. M. Hare,^a M. J. Binks,^a K. Grimwood,^{b,c} A. B. Chang,^{a,b,d,e} A. J. Leach,^a and H. Smith-Vaughan^a

Child Health Division, Menzies School of Health Research, Charles Darwin University, Darwin, Northern Territory,^a Queensland Children's Medical Research Institute, The University of Queensland, Royal Children's Hospital, Brisbane, Queensland,^b Department of Infectious Diseases, Royal Children's Hospital, Brisbane, Queensland,^c Royal Darwin Hospital, Darwin, Northern Territory,^d and Department of Respiratory Medicine, Royal Children's Hospital, Brisbane, Queensland,^e Australia

A PCR for protein D (hpd#3) was used to differentiate nontypeable *Haemophilus influenzae* (NTHI) from *Haemophilus haemolyticus*. While 90% of nasopharyngeal specimens and 100% of lower-airway specimens from 84 Indigenous Australian children with bronchiectasis had phenotypic NTHI isolates confirmed as *H. influenzae*, only 39% of oropharyngeal specimens with phenotypic NTHI had *H. influenzae*. The nasopharynx is therefore the preferred site for NTHI colonization studies, and NTHI is confirmed as an important lower-airway pathogen.

Nontypeable *Haemophilus influenzae* (NTHI) colonizes the upper airways, where it is an important cause of otitis media (10). It is also isolated frequently from the lower airways of adults with chronic obstructive pulmonary disease (COPD) (8) and children and adults with bronchiectasis (3, 4).

Accurate identification of NTHI is important, since nonhemolytic strains of the closely related (primarily commensal) *Haemophilus haemolyticus* may be misidentified as NTHI by phenotypic methods used in most clinical microbiology laboratories (9). Molecular detection techniques have revealed that 12 to 27% of nasopharyngeal isolates from healthy and otitis-prone children, which were initially identified by classical phenotypic methods as NTHI isolates (here referred to as phenotypic NTHI isolates), were actually *H. haemolyticus* (5, 9), while *H. haemolyticus* was not identified in middle ear fluid samples from children with acute otitis media (5). In contrast, *H. haemolyticus* comprised 40% of phenotypic NTHI sputum isolates from adults with COPD (9).

These findings raise important questions about NTHI as a true lower respiratory pathogen. We therefore investigated the proportion of *H. haemolyticus* isolates among phenotypic NTHI isolates in the upper and lower airways of Indigenous Australian children with bronchiectasis, a population with high rates of NTHI colonization and associated respiratory disease (2, 3, 12).

The Human Research Ethics Committee of the Northern Territory Department of Health and Menzies School of Health Research approved the study. Following written, informed parent/guardian consent, paired nasopharyngeal swabs and bronchoalveolar lavage (BAL) fluid specimens were collected from a convenience sample of 84 (62% male) Indigenous children aged 5 to 155 (median, 27) months who were undergoing routine diagnostic evaluation following radiographic confirmation of bronchiectasis at the Royal Darwin Hospital between July 2007 and December 2010. Oropharyngeal swabs were also collected from 56 of these children. Specimens were stored and cultured using standard microbiologic methods (3). Lower-airway infection was defined by a semiquantitative growth score of ≥ 4 , which correlated with $>10^4$ CFU/ml BAL fluid, as determined by serial dilution and quantitative counts (3).

Phenotypic NTHI isolates were identified by morphology, re-

quirement for X and V growth factors, and failure to react with Phadebact (Bactus AB, Sweden) antisera specific for *H. influenzae* type b or types a and c through f. Up to 4 colonies (including any with differing morphology) were isolated from each culture-positive specimen and tested using PCR. Confirmation of phenotypic NTHI isolates was performed using defined DNA extraction methods (12) and a TaqMan-based, real-time PCR assay targeting protein D (hpd#3), which discriminates between *H. influenzae* and *H. haemolyticus* isolates (1, 13). Phenotypic NTHI isolates returning negative PCR results were considered to be *H. haemolyticus*, since the only other X- and V-factor-dependent *Haemophilus* species (*H. aegyptius*, an important cause of conjunctivitis) has a different appearance, requires additional growth factors, and is unlikely to be cultured from these sites (6).

Table 1 shows the numbers, proportions, and distribution of phenotypic NTHI, hpd#3 PCR-confirmed *H. influenzae*, presumptive *H. haemolyticus*, and concurrent *H. influenzae* and *H. haemolyticus* in the 224 specimens collected from the upper and lower airways. A total of 214 isolates from 108 phenotypic NTHI-positive specimens (an average of 1.8 colonies isolated per nasopharyngeal and oropharyngeal swab and 2.4 colonies per BAL specimen) were tested. Most nasopharyngeal (87%) and BAL fluid (88%) isolates were confirmed as *H. influenzae*, but almost two-thirds (65%) of oropharyngeal isolates were presumptive *H. haemolyticus*. *H. influenzae* and *H. haemolyticus* were isolated concurrently from 10% (4/42) of nasopharyngeal swabs, 17% (6/36) of oropharyngeal swabs, and 27% (8/30) of BAL fluid cultures from children with phenotypic NTHI carriage or lower-airway infection.

This study shows that, similar to healthy and otitis-prone Western

Received 1 March 2012 Returned for modification 27 March 2012

Accepted 20 April 2012

Published ahead of print 2 May 2012

Address correspondence to K. M. Hare, kim.hare@menzies.edu.au.

Copyright © 2012, American Society for Microbiology. All Rights Reserved.

doi:10.1128/JCM.00566-12

TABLE 1 Culture and PCR results for phenotypic NTHI isolates from nasopharyngeal and oropharyngeal swabs and bronchoalveolar lavage fluid specimens from 84 Australian Indigenous children with bronchiectasis^a

Isolate source	No. of specimens	No. (%) of specimens with:			
		Phenotypic NTHI	Phenotypic NTHI confirmed by PCR as <i>H. influenzae</i>	Presumptive <i>H. haemolyticus</i>	Concurrent <i>H. influenzae</i> and presumptive <i>H. haemolyticus</i>
NP swab	84	42 (50)	38 (45)	8 (10)	4 (5)
OP swab	56	36 (64)	14 (25)	28 (50)	6 (11)
BAL fluid	84	30 ^b (36)	30 (36)	8 (10)	8 (10)
Total	224	108 (48)	82 (37)	44 (20)	18 (8)

^a NP, nasopharyngeal; OP, oropharyngeal; BAL, bronchoalveolar lavage; NTHI, nontypeable *Haemophilus influenzae*.

^b Specimens with phenotypic NTHI lower-airway infection (>10⁴ CFU/ml BAL fluid).

Australian children (5), most phenotypic NTHI nasopharyngeal isolates from Indigenous children with bronchiectasis were confirmed as *H. influenzae*. In contrast, many apparent NTHI isolates from oropharyngeal swabs were *H. haemolyticus* in our study population. Previous oropharyngeal culture-based studies of *H. influenzae* may have therefore overestimated NTHI carriage, and our data instead support the nasopharynx as the preferred site for *H. influenzae* carriage studies in children. Such studies are important to monitor antimicrobial resistance and detect changes in pharyngeal biota that may result from antibiotic administration or vaccination.

Quantifying pathogens in BAL fluid helps adjust for upper-airway contamination during bronchoscopy. Our finding that 100% of BAL specimens with phenotypic NTHI lower-airway infection (>10⁴ CFU/ml BAL fluid) were PCR positive for *H. influenzae* confirms NTHI as a lower-airway pathogen. However, the role of *H. haemolyticus* is unclear. Prior studies report that *H. haemolyticus* is rarely found in sterile sites, including middle ear fluid, or associated with clinically defined infections (5, 7, 9, 11). While our findings suggest that *H. haemolyticus* has a propensity for the oropharynx, without specific molecular detection and quantification, we cannot determine whether its presence in lower-airway cultures represents upper-airway contamination or a pathogenic role.

In conclusion, we have used *H. influenzae*-specific PCR to reaffirm the importance of NTHI as a lower-airway pathogen in Australian Indigenous children with bronchiectasis. In addition, we have shown that the nasopharynx, rather than the oropharynx, is the preferred site for NTHI carriage studies in this population.

ACKNOWLEDGMENTS

We thank the children and their caregivers for participating in the study, Menzies staff (Gabrielle MacCallum, Kobi Schutz, and Susan Pizzutto) and Royal Darwin Hospital staff for assistance with specimen collection, Menzies laboratory staff (Vanya Hampton, Joanna Bugg, Estelle Carter,

and Peter Christensen) for assistance with laboratory processing, and Robyn Marsh for helpful comments.

This study was supported by NHMRC project grant 545223.

REFERENCES

- Binks MJ, et al. 2012. Molecular surveillance of true nontypeable *Haemophilus influenzae*: an evaluation of PCR screening assays. PLoS One 7:e34083. doi:10.1371/journal.pone.0034083.
- Chang AB, Masel JP, Boyce NC, Wheaton G, Torzillo PJ. 2003. Non-CF bronchiectasis: clinical and HRCT evaluation. Pediatr. Pulmonol. 35:477–483.
- Hare KM, et al. 2010. Respiratory bacterial pathogens in the nasopharynx and lower airways of Australian indigenous children with bronchiectasis. J. Pediatr. 157:1001–1005.
- King PT, Holdsworth SR, Freezer NJ, Villanueva E, Holmes PW. 2007. Microbiologic follow-up study in adult bronchiectasis. Respir. Med. 101:1633–1638.
- Kirkham LA, et al. 2010. Nasopharyngeal carriage of *Haemophilus haemolyticus* in otitis-prone and healthy children. J. Clin. Microbiol. 48:2557–2559.
- Ledeboer NA, Doern GV. 2012. *Haemophilus*, p 588–602. In Versalovic J, et al (ed), Manual of clinical microbiology, 10th ed. ASM Press, Washington, DC.
- Morton DJ, Hempel RJ, Whitby PW, Seale TW, Stull TL. 2012. An invasive *Haemophilus haemolyticus* isolate. J. Clin. Microbiol. 50:1502–1503.
- Murphy TF. 2003. Respiratory infections caused by non-typeable *Haemophilus influenzae*. Curr. Opin. Infect. Dis. 16:129–134.
- Murphy TF, et al. 2007. *Haemophilus haemolyticus*: a human respiratory tract commensal to be distinguished from *Haemophilus influenzae*. J. Infect. Dis. 195:81–89.
- Murphy TF, et al. 2009. Nontypeable *Haemophilus influenzae* as a pathogen in children. Pediatr. Infect. Dis. J. 28:43–48.
- Norskov-Lauritsen N. 2009. Detection of cryptic genospecies misidentified as *Haemophilus influenzae* in routine clinical samples by assessment of marker genes *fucK*, *hap*, and *sodC*. J. Clin. Microbiol. 47:2590–2592.
- Smith-Vaughan H, et al. 2006. Measuring nasal bacterial load and its association with otitis media. BMC Ear Nose Throat Disord. 6:10.
- Wang X, et al. 2011. Detection of bacterial pathogens in Mongolia meningitis surveillance with a new real-time PCR assay to detect *Haemophilus influenzae*. Int. J. Med. Microbiol. 301:303–309.