

COMMONWEALTH OF AUSTRALIA

Copyright Regulations 1969

Warning

This material has been reproduced and communicated to you by or on behalf of *The Charles Darwin University* pursuant to Part VB of the *Copyright Act 1968* (the Act). The material in this communication may be subject to copyright under the Act. Any further reproduction or communication of this material by you may be the subject of copyright protection under the Act.

Do not remove this notice



Family Name	
Given Names	
Student Number	
Teaching Period	Semester 2, 2016

FINAL EXAMINATION	DURATION
TEP024 – Advanced Foundation Maths	Reading Time: 10 minutes
	Writing Time: 180 minutes

INSTRUCTIONS TO CANDIDATES

- 1.1 All questions are to be answered in the examination booklet provided
- 1.2 Read ALL questions carefully and attempt to answer all 15 questions.
- 1.3 Full working on all questions must be shown.
- 1.4 The maximum mark that can be obtained on the exam is 120 Marks.

EXAM CONDITIONS

You may begin writing from the commencement of the examination session.

The reading time indicated above is provided as a guide only.

This is a RESTRICTED OPEN BOOK examination

Any calculator is permitted

One A4 sheet of handwritten double-sided notes permitted

Any hard copy, unannotated English dictionary is permitted

ADDITIONAL AUTHORISED MATERIALS	EXAMINATION MATERIALS TO BE SUPPLIED
No additional printed material is permitted	2 x Scrap Paper

**THIS EXAMINATION IS PRINTED
DOUBLE-SIDED.**

THIS PAGE HAS BEEN INTENTIONALLY LEFT

Question 1.**[10 Marks]**

- (a) Simplify and rewrite the following algebraic expression using positive indices only. [3]

$$\frac{(3x^2)^3}{x^5y} \div \frac{(-3y^4)^2}{x}$$

- (b) Solve for the unknown x in the following algebraic equation. [3]

$$3(2x - 7) = -2(x - 7) - 3$$

- (c) Two points with coordinates $(0,1)$ and $(-3,-14)$ are joined to form a straight line. [4]

(i) Determine the gradient of this line.

(ii) Write the equation of the line.

Question 2.**[5 Marks]**

- (a) Given the equation $F = \frac{GMm}{r^2}$ evaluate to find F , if
 $G = 6.67 \times 10^{-11}$, $M = 5.98 \times 10^{24}$, $r = 1000$, $m = 56$.

Give your answer in scientific notation correct to 3 significant figures. [3]

- (b) Transpose the formulae $F = \frac{GMm}{r^2}$ to make r the subject of the formula. [2]

Question 3.**[4 Marks]**

Solve the following system of linear equations for x and y algebraically giving your answers as rational fractions. (Use the Elimination method for both x and y).

$$6x - y = 5$$

$$-\frac{1}{3}x - y = -2$$


Question 4.**[13 Marks]**

(a) Given the function $f(x) = (3 - x)(x + 5)$, expand the expression and hence rewrite in expanded quadratic form. [2]

(b) (i) State the coordinates of the X and Y intercepts of the function. [3]

(ii) Find the coordinates of the turning point of the function. [2]

(c) Sketch the function on the axes given below **labelling** all axis intercepts (X and Y intercepts) with their **coordinates**, and **label** the turning point with its **coordinates**. [4]



(d) Find by any means possible (algebraically or otherwise) the solution for $f(x) = (3 - x)(x + 5) = -9$. [2]

Question 5.**[5 Marks]**

Using the quadratic formula, solve $2x^2 - 6x + 3 = 0$ for all values of x . Give your answer in simplest surd form. (State the values of a , b and c before you attempt to use the quadratic formula.)

Question 6.**[3 Marks]**

Solve the following equation for x without the use of logarithms.

$$2^{1-2x} = \frac{1}{32}$$

Question 7.**[8 Marks]**

(a) Rewrite $3\log 2 - \log 5 + 2$ as a single logarithm.

[3]

(b) Simplify $\frac{8 \log x}{\log x^2}$ without the use of a calculator.

[2]

(c) $\log 5 + 3\log x^2 - 2\log x$

[3]

Question 8.**[7 Marks]**Solve the following equations for t (correct to 3 sig. figs.).

(a) $4e^{0.36t} = 100$ [3]

(b) $2^t = 5^{1-t}$ [4]

Question 9.**[6 Marks]**

Two towns on the earth's equator are separated by 1.5° in longitude


- (a) Convert 1.5° to radians and give your answer correct to 4 significant figures. [2]
- (b) Given the earth's radius is $6.38 \times 10^6 \text{ m}$ and $\left(\frac{1}{60}\right)^\circ$ represents 1852 m on the ground at the equator, what is the distance between the two towns? [2]
- (c) Determine the speed (in km/h) of a vehicle which can traverse the distance between the two towns in 2 hours and 15 minutes. [2]

Question 10.**[9 Marks]**

A vertical mobile phone tower OM has a height of 37.8 m. A stainless steel supporting cable OA makes an angle of 63° with the ground. A second cable OB is 63.4 m in length.

Give all answers correct to 3 sig. figs.

- (a) Use your knowledge of right angle trigonometry to find the length of the cable OA. [3]



- (b) Demonstrate your knowledge of non-right angle trigonometry (cosine or sine rules) to find the angle between the ground and the second cable ($\angle ABO$), and the distance AB between the two cables on the ground. [6]

Question 11.**[8 Marks]**


(a) If $\sin\beta = -0.687$ and $0^\circ \leq \beta \leq 360^\circ$ find β . [3]

(b) Use your knowledge of the laws of logarithms to solve $150e^{-0.26t} = 12(1.28)^t$
(Give answer correct to 4 sig. figs.) [5]

Question 12.**[7 Marks]**Given the function $f(x) = -3\sin 2x$

- (a) State the value of the amplitude and the period of the function. [2]


- (b) Sketch the graph of the function on the axes given below. [2]



- (c) Determine algebraically all solutions to $f(x) = 2$ that can be demonstrated graphically on the graph that you have drawn above. (ie all solutions for $f(x) = 2$ given the domain $0 \leq x \leq 270^\circ$) [3]

Question 13.**[10 marks]**

- (a) Two velocity vectors 50.0ms^{-1} 053° and 72.0ms^{-1} 295° are added. Complete the diagram below which is to scale, and hence determine the magnitude and direction of the resultant velocity by measurement. SCALE: $1\text{cm} : 10\text{ms}^{-1}$. [3]



- (b) Use your knowledge of trigonometry to verify your answer in part (a) above (correct to 3 sig. figs.). [7]

Question 14.

[15 Marks]


Drawn below is the graph of the function $N(t) = 170(0.965)^t$ where N represents the number of termites in a laboratory termite mound over time, t , in weeks. These termites are given little food and poor living conditions.



- (a) (i) Calculate the initial value of N , ie $N(0)$. [1]
- (ii) Plot this point on your graph labelling the point with its coordinates. [1]
- (b) (i) Calculate the number of termites after one year. [1]
- (ii) Plot this point on the graph labelling it with its coordinates. [1]
- (c) Determine algebraically how many weeks it will take for the number of termites in the mound to decrease to 25% of the original number. [4]

this question is continued on the next page.

- (d) Another mound of termites in the same laboratory is given the choice slivers of bark and wood to eat, and ideal living conditions. Their numbers are found to increase over time (t , in weeks) according to the model $P(t) = 50(1.026)^t$. Plot the graph for this model on the axes given below. Show at least 3 points along the curve labelling each point with its coordinates. [3]



- (e) At some point in time the number of termites in both colonies will be equal. Determine when that will be to the nearest whole week and how many termites there will be in each colony at that time.

ie Solve $170(0.965)^t = 50(1.026)^t$ [4]

Question 15.

[10 Marks]

- (a) Two force vectors $P = 48N \ 316^\circ$ and $Q = 123N \ 220^\circ$ in the rectangular coordinate system are added. Draw an accurate scale diagram so that you can measure the magnitude and direction of the addition of these two vectors. [3]

this question is continued on the next page.

- (b) Add the two vectors by finding their horizontal and vertical components and hence calculate the magnitude and direction of the resultant vector. (Work correct to 3 decimal places and give final answer correct to the nearest whole Newton (N) and whole degree ($^\circ$)). [7]

		R_x	R_y
A	$P = 48N \ 316^\circ$		
B	$Q = 123N \ 220^\circ$		

MAGNITUDE

DIRECTION

RESULTANT